arXiv:1512.03255v1 [math.NT] 10 Dec 2015

ON THE CLASS OF $2D$ q-APPELL POLYNOMIALS

MARZIEH EINI KELESHTERI AND NAZIM I. MAHMUDOV

Abstract. In this research, as the new results of our previously proposed definition for the new class of $2D$ q-Appell polynomials in [1], we derive some interesting relations including the recurrence relation and partial q-difference equation of the aforementioned family of q-polynomials. Next, as some famous examples of this new defined class of q-polynomials, we obtain the corresponding relations to the $2D$ q-Bernoulli polynomials, $2D$ q-Euler polynomials as well as $2D$ q-Genocchi polynomials.

1. Introduction

In [1], Eini and Mahmudov defined $2D$ q-Appell Polynomials by means of the following generating function

$$A_q(x,y;t) := A_q(t)e_q(tx)E_q(ty) = \sum_{n=0}^{\infty} A_{n,q}(x,y)\frac{t^n}{[n]_q!}, \tag{1}$$

where

$$A_q(t) := \sum_{n=0}^{\infty} A_{n,q}\frac{t^n}{[n]_q!}, \quad A_q(t) \neq 0, \tag{2}$$

is an analytic function at $t = 0$, and $A_{n,q} := A_{n,q}(0,0)$. Taking q-derivative of $A_q(x,y,t)$ with respect to the variable x, from one hand we obtain

$$\begin{aligned} D_{q,x}(A_q(x,y;t)) &= D_{q,x}(A_q(t)e_q(tx)E_q(ty)) \\ &= tA_q(t)e_q(tx)E_q(ty) = \sum_{n=0}^{\infty} A_{n,q}(x,y)\frac{t^{n+1}}{[n]_q!} \\ &= \sum_{n=1}^{\infty} [n]_q\, A_{n-1,q}(x,y)\frac{t^n}{[n]_q!}. \end{aligned} \tag{3}$$

1991 *Mathematics Subject Classification.* Special Functions.

Key words and phrases. q-polynomials, lowering operator, $2D$, Appell, Euler, Bernoulli, Genocchi.

From another hand we can write

$$D_{q,x}(A_q(x,y;t)) = D_{q,x}(\sum_{n=0}^{\infty} A_{n,q}(x,y)\frac{t^n}{[n]_q!})$$

$$= \sum_{n=0}^{\infty} D_{q,x}(A_{n,q}(x,y))\frac{t^n}{[n]_q!}. \tag{4}$$

Comparing the coefficients of $\frac{t^n}{[n]_q!}$ in the relations (3) and (4), leads to obtain

$$D_{q,x}(A_{n,q}(x,y)) = [n]_q\, A_{n-1,q}(x,y). \tag{5}$$

Using a similar technique for taking q-derivative of $A_q(x,y;t)$ with respect to the variable y, we have

$$D_{q,y}(A_{n,q}(x,y)) = [n]_q\, A_{n-1,q}(x,qy). \tag{6}$$

Now, according to relations (5) and (6), we define the following lowering operators

$$\Phi_{n,q_x} = \frac{1}{[n]_q} D_{q,x}, \quad \Phi_{n,q_y} = \frac{1}{[n]_q} D_{q,y}. \tag{7}$$

Therefore, we may reexpress the relations (5) and (6)in the form of the following operational identities

$$\Phi_{n,q_x} A_{n,q}(x,y) = A_{n-1,q}(x,y), \text{ and } \quad \Phi_{n,q_y} A_{n,q}(x,y) = A_{n-1,q}(x,qy), \tag{8}$$

respectively. Eventually applying the above operators k times, leads to obtain

$$A_{n-k,q}(x,y) = \big(\Phi_{n-k,q_x} \circ \ldots \circ \Phi_{n,q_x}\big) A_{n,q}(x,y) = \frac{[n-k]_q!}{[n]_q!} D^k_{q,x} A_{n,q}(x,y), \tag{9}$$

and

$$A_{n-k,q}(x,q^k y) = \big(\Phi_{n-k,q_y} \circ \ldots \circ \Phi_{n,q_y}\big) A_{n,q}(x,y) = \frac{[n-k]_q!}{[n]_q!} D^k_{q,y} A_{n,q}(x,y), \tag{10}$$

respectively.

2. Recurrence Relations And q-Difference Equations For The Class Of $2D$ q-Appell Polynomials

In 2002, Bretti et. al. proposed a generating function for the family of $2D$ Appell polynomials, [2]. They, also, obtained the corresponding recurrence relations and differential equations to the aforementioned family by calculating raising and lowering operators. In [3], Mahmudov applied an innovative technique in order to derive the recurrence relations and difference equations of the polynomials in the class of q-Appell polynomials only by using only lowering operators that are q-derivatives. Inspired by his novel approach, in the following we will use a similar technique in order to derive the corresponding relations to the class of $2D$ q-Appell polynomials.

Theorem 1. *The following linear homogeneous recurrence relation holds for the class of $2D$ q-Appell polynomials*

(11)
$$A_{n,q}(qx,y)=\frac{1}{[n]_q}\sum_{k=1}^{n}\left[\begin{array}{c} n \\ k \end{array}\right]_q q^{n-k}A_{n-k}(x,y)(\alpha_k+\frac{\beta_{k-1}}{[k]_q}y)+xq^nA_{n-1,q}(x,y),\quad n\geq 1,$$

or equivalently,

(12)
$$A_{n,q}(qx,y)=q^n(x+(\alpha_1+\beta_0y)q^{-1})A_{n-1,q}(x,y)+$$
$$\frac{1}{[n]_q}\sum_{k=1}^{n-1}\left[\begin{array}{c} n \\ k-1 \end{array}\right]_q q^{k-1}A_{k-1,q}(x,y)(\alpha_{n-k+1}+\frac{y}{[n-k+1]_q}\beta_{n-k}),\quad n\geq 1.$$

Proof. Starting with taking the q-derivative of the generating function in relation (1) with respect to t, we have

$$D_{q,t}(A_q(qx,y;t))=$$

(13)
$$yA_q(t)e_q(qtx)E_q(qty)+D_{q,t}(A_q(t))e_q(qtx)E_q(qty)+qxA_q(x,y;qt).$$

Now, multiplying both sides of the identity (13) by t and factorizing $A_q(x,y;qt)$ form its left hand side, we obtain

$$tD_{q,t}(A_q(qx,y;t)) = A_q(x,y;qt)\Big[t\frac{D_{q,t}(A_q(t))}{A_q(qt)} + tqx + ty\frac{A_q(t)}{A_q(tq)}\Big]. \tag{14}$$

Suppose that $t\frac{D_{q,t}(A_q(t))}{A_q(qt)} = \sum\limits_{n=0}^{\infty}\alpha_n\frac{t^n}{[n]_q!}$, and $t\frac{A_q(t)}{A_q(qt)} = \sum\limits_{n=0}^{\infty}\beta_n\frac{t^n}{[n]_q!}$. Starting from taking q-derivative of the left hand side of relation (14) with respect to t and also substituting the assumptions above in the right hand side of the same equation, we can continue as

$$\sum_{n=1}^{\infty}[n]_q A_{n,q}(qx,y)\frac{t^n}{[n]_q!} = A_q(x,y;qt)\Big[\sum_{n=0}^{\infty}\alpha_n\frac{t^n}{[n]_q!} + \sum_{n=0}^{\infty}y\beta_n\frac{t^{n+1}}{[n]_q!} + tqx\Big]. \tag{15}$$

The last part of identity above can be written as

$$= \sum_{n=0}^{\infty}q^n A_{n,q}(x,y)\frac{t^n}{[n]_q!}\Big[\sum_{n=0}^{\infty}\alpha_n\frac{t^n}{[n]_q!} + y\sum_{n=1}^{\infty}[n]_q\beta_n\frac{t^n}{[n]_q!} + tqx\Big], \tag{16}$$

which is equivalent to

$$= \sum_{n=0}^{\infty}\sum_{k=0}^{n}\Big(\begin{bmatrix} n \\ k \end{bmatrix}_q q^{n-k}A_{n-k,q}(x,y)\alpha_k\Big)\frac{t^n}{[n]_q!} + y\sum_{n=0}^{\infty}\sum_{k=0}^{n}\Big(\begin{bmatrix} n \\ k \end{bmatrix}_q q^{n-k}A_{n-k,q}(x,y)\beta_k\Big)\frac{t^{n+1}}{[n]_q!} + x\sum_{n=0}^{\infty}q^{n+1}A_{n,q}(x,y)\frac{t^{n+1}}{[n]_q!}.$$

This means that

$$\sum_{n=1}^{\infty}[n]_q\,A_{n,q}(qx,y)\frac{t^n}{[n]_q!}=A_{0,q}(x,y)\alpha_0+$$

$$\sum_{n=1}^{\infty}\left(\sum_{k=0}^{n}\left[\begin{array}{c}n\\k\end{array}\right]_q q^{n-k}A_{n-k,q}(x,y)\alpha_k+y\sum_{k=0}^{n-1}\left[\begin{array}{c}n-1\\k\end{array}\right]_q[n]_q q^{n-k-1}A_{n-k-1,q}(x,y)\beta_k+\right.$$

(17)

$$\left.x[n]_q^n A_{n-1,q}(x,y)\right)\frac{t^n}{[n]_q!}.$$

$$(18)\quad =\sum_{n=0}^{\infty}\sum_{k=0}^{n}\Big(\left[\begin{array}{c}n\\k\end{array}\right]_q q^{n-k}A_{n-k}(x,y)(\alpha_k+\beta_k y)+[n]_q x q^n A_{n-1,q}(x,y)\Big)\frac{t^n}{[n]_q!}.$$

Comparing the coefficients of $\frac{t^n}{[n]_q!}$ in both sides of relation (17) and noting to the fact that $\alpha_0=0$, lead to obtain the following identity for $n\geq 1$

$$[n]_q A_{n,q}(qx,y)=\sum_{k=1}^{n}\left[\begin{array}{c}n\\k\end{array}\right]_q q^{n-k}A_{n-k,q}(x,y)\alpha_k+$$

$$y\sum_{k=1}^{n}\left[\begin{array}{c}n-1\\k-1\end{array}\right]_q[n]_q q^{n-k}A_{n-k,q}(x,y)\beta_{k-1}+x[n]_q q^n A_{n-1,q}(x,y),$$

whence the result. □

Theorem 2. *The following partial q-difference equations hold for the polynomials in the class of 2D q-Appell*

$$(19)\quad \left[\sum_{k=1}^{n}\frac{q^{n-k}}{[k]_q!}(\alpha_k+\frac{\beta_{k-1}}{[k]_q}y)D_{q,x}^k+xq^nD_{q,x}\right]A_{n,q}(x,y)-[n]_qA_{n,q}(qx,y)=0.$$

(20)

$$\sum_{k=1}^{n}\frac{q^{n-k}}{[k]_q!}(\alpha_k+\frac{\beta_{k-1}}{q^k[k]_q}y)D_{q,y}^kA_{n,q}(x,\frac{y}{q^k})+xq^nD_{q,y}A_{n,q}(x,\frac{y}{q})-[n]_qA_{n,q}(qx,y)=0.$$

Proof. The proof is the direct result of replacing relations (9) and (10) in the linear homogeneous recurrence relation (11) given in Theorem (1), respectively. □

3. q-Difference Equations For Various Members Of The Family Of q-Appell Polynomials

Choosing different functions as $A_q(t)$ in Definition (1), leads to generate different members of $2D$ q-Appell polynomials. In the following we introduce some of the most famous $2D$ q-Appell polynomials and derive the corresponding recurrence relations and partial q-difference equations to them.

3.1. $2D$ q-Bernoulli polynomials.

Taking $A_q(t) = \frac{t}{e_q(t)-1}$ in Definition (1), leads to obtain $2D$ q-Bernoulli polynomials, $B_{n,q}(x,y)$, [4], [5].

$$B_q(x,y;t) := \frac{t}{e_q(t)-1} e_q(tx) E_q(ty) = \sum_{n=0}^{\infty} B_{n,q}(x,y) \frac{t^n}{[n]_q!}, \tag{21}$$

Lemma 3. *Suppose that*

$$t\frac{D_{q,t}(A_q(t))}{A_q(qt)} = t\frac{D_{q,t}\frac{t}{e_q(t)-1}}{\frac{qt}{e_q(qt)-1}} = \sum_{n=0}^{\infty} \alpha_n \frac{t^n}{[n]_q!}, \tag{22}$$

and

$$\frac{A_q(t)}{A_q(qt)} = \frac{\frac{t}{e_q(t)-1}}{\frac{qt}{e_q(qt)-1}} = \sum_{n=0}^{\infty} \beta_n \frac{t^n}{[n]_q!}, \tag{23}$$

then

$$\alpha_n = \frac{-1}{q} b_{n,q}, \quad \alpha_1 = \frac{-1}{[2]_q}, \tag{24}$$

and

$$\beta_n = \frac{q-1}{q} \sum_{k=0}^{n} \begin{bmatrix} n \\ k \end{bmatrix}_q b_{k,q}, \quad \text{for } n \geq 1 \quad \text{and } \beta_0 = 1, \tag{25}$$

where $b_{n,q} = B_{n,q}(0,0)$ is the n-th q-Bernoulli number and can be obtained from the generating function $\frac{t}{e_q(t)-1} = \sum_{n=0}^{\infty} b_{n,q} \frac{t^n}{[n]_q!}$.

Theorem 4. *The following linear homogeneous recurrence relation holds for the class of 2D q-Bernoulli polynomials for every* $n \geq 1$

$$B_{n,q}(qx,y) = q^n(x + (\frac{-1}{[2]_q} + y)q^{-1})B_{n-1,q}(x,y)+$$

(26)

$$\frac{1}{[n]_q}\sum_{k=1}^{n-1}\left[\begin{array}{c} n \\ k-1 \end{array}\right]_q q^{k-2}B_{k-1,q}(x,y)(-b_{n-k+1,q} + \frac{(q-1)y}{[n-k+1]_q}\sum_{l=0}^{n-k}\left[\begin{array}{c} n-k \\ l \end{array}\right]_q b_{l,q}).$$

Theorem 5. *The following partial q-difference equations hold for the polynomials in the class of 2D q-Bernoulli*

$$\left[(xq^n + y - \frac{1}{[2]_q})D_{q,x} + \sum_{k=2}^{n}\frac{q^{n-k-1}}{[k]_q!}(-b_{k,q} + (q-1)y\sum_{l=0}^{k-1}\left[\begin{array}{c} k-1 \\ l \end{array}\right]_q b_{l,q})D_{q,x}^k\right]$$

(27)

$$\times B_{n,q}(x,y) - [n]_q B_{n,q}(qx,y) = 0.$$

$$(xq^n + \frac{y}{q} - \frac{1}{[2]_q})D_{q,y}A_{n,q}(x,\frac{y}{q}) + \sum_{k=2}^{n}\frac{q^{n-k-1}}{[k]_q!}(-b_{k,q} + \frac{(q-1)y}{q^k[k]_q}\sum_{l=0}^{k}\left[\begin{array}{c} k \\ l \end{array}\right]_q b_{l,q})$$

(28)

$$\times D_{q,y}^k B_{n,q}(x,\frac{y}{q^k}) - [n]_q A_{n,q}(qx,y) = 0.$$

3.2. $2D$ q**-Euler polynomials.** Taking $A_q(t) = \frac{2}{e_q(t)+1}$ in Definition (1), leads to obtain $2D$ q-Euler polynomials, $E_{n,q}(x,y)$, as follows[4], [5].

$$E_q(x,y;t) := \frac{2}{e_q(t)+1}e_q(tx)E_q(ty) = \sum_{n=0}^{\infty}E_{n,q}(x,y)\frac{t^n}{[n]_q!}, \tag{29}$$

Lemma 6. *Suppose that*

$$t\frac{D_{q,t}(A_q(t))}{A_q(qt)} = t\frac{D_{q,t}\frac{2}{e_q(t)+1}}{\frac{2}{e_q(qt)+1}} = \sum_{n=0}^{\infty}\alpha_n\frac{t^n}{[n]_q!}, \tag{30}$$

and

$$\frac{A_q(t)}{A_q(qt)} = \frac{\frac{2}{e_q(t)+1}}{\frac{2}{e_q(qt)+1}} = \sum_{n=0}^{\infty}\beta_n\frac{t^n}{[n]_q!}, \tag{31}$$

then

$$\alpha_n = \frac{1}{2}E_{n-1,q}, \quad \alpha_1 = \frac{-1}{2}, \tag{32}$$

and

$$\beta_n = \frac{q-1}{2}\sum_{k=0}^{n}\begin{bmatrix} n \\ k \end{bmatrix}_q E_{k,q}, \quad \text{for } n \geq 1 \quad \text{and } \beta_0 = \frac{q+1}{2}, \tag{33}$$

where $E_{n,q} = E_{n,q}(0,0)$ *is the n-th q-Euler number and can be obtained from the generating function* $\frac{2}{e_q(t)+1} = \sum_{n=0}^{\infty} E_{n,q}\frac{t^n}{[n]_q!}$.

Theorem 7. *The following linear homogeneous recurrence relation holds for the class of* $2D$ *q-Euler polynomials for every* $n \geq 1$

$$A_{n,q}(qx,y) = q^n(x + \frac{(q+1)y-1}{2q})E_{n-1,q}(x,y)+$$
$$\frac{1}{[n]_q}\sum_{k=1}^{n-1}\begin{bmatrix} n \\ k-1 \end{bmatrix}_q q^{k-1}E_{k-1,q}(x,y)(\frac{1}{2}E_{n-k,q}+\frac{(q-1)y}{q[n-k+1]_q}\sum_{l=0}^{n-k}\begin{bmatrix} n-k \\ l \end{bmatrix}_q E_{l,q}).$$

Theorem 8. *The following partial q-difference equations hold for the polynomials in the class of* $2D$ *q-Euler*

$$\left[(xq^n + \frac{(q+1)y}{2[k]_q} - \frac{1}{2})D_{q,x} + \sum_{k=2}^{n}\frac{q^{n-k}}{[k]_q!}(\frac{1}{2}E_{k-1,q} + \frac{(q-1)y}{2[k]_q}\sum_{l=0}^{k-1}\begin{bmatrix} k-1 \\ l \end{bmatrix}_q E_{l,q})D_{q,x}^k\right]$$
$$\times E_{n,q}(x,y) - [n]_q E_{n,q}(qx,y) = 0.$$

$$(xq^n + \frac{(q+1)y}{2q[k]_q} - \frac{1}{2})D_{q,y}E_{n,q}(x,\frac{y}{q}) + \sum_{k=2}^{n}\frac{q^{n-k}}{[k]_q!}(\frac{1}{2}E_{k-1,q} + \frac{(q-1)y}{2q^k[k]_q}\sum_{l=0}^{k-1}\begin{bmatrix} k-1 \\ l \end{bmatrix}_q E_{l,q})$$
$$\times D_{q,y}^k E_{n,q}(x,\frac{y}{q^k}) - [n]_q E_{n,q}(qx,y) = 0.$$

3.3. $2D$ **q-Genocchi polynomials.** Taking $A_q(t) = \frac{2t}{e_q(t)+1}$ in Definition (1), leads to obtain $2D$ q-Genocchi polynomials, $G_{n,q}(x,y)$, as follows

$$G_q(x,y;t) := \frac{2t}{e_q(t)+1}e_q(tx)G_q(ty) = \sum_{n=0}^{\infty}G_{n,q}(x,y)\frac{t^n}{[n]_q!}, \tag{34}$$

Lemma 9. *Suppose that*

$$t\frac{D_{q,t}(A_q(t))}{A_q(qt)} = t\frac{D_{q,t}\frac{2t}{e_q(t)+1}}{\frac{2tq}{e_q(qt)+1}} = \sum_{n=0}^{\infty}\alpha_n\frac{t^n}{[n]_q!}, \tag{35}$$

and

$$\frac{A_q(t)}{A_q(qt)} = \frac{\frac{2t}{e_q(t)+1}}{\frac{2tq}{e_q(qt)+1}} = \sum_{n=0}^{\infty}\beta_n\frac{t^n}{[n]_q!}, \tag{36}$$

then

$$\alpha_n = \frac{1}{2q}G_{n,q}, \quad \textit{for } n\geq 2, \textit{ and } \alpha_0 = \frac{1}{q},\ \alpha_1 = \frac{-1}{q}, \tag{37}$$

and

$$\beta_n = \frac{q-1}{2q}\sum_{k=0}^{n}\begin{bmatrix} n \\ k \end{bmatrix}_q G_{k,q}, \quad \textit{for } n\geq 1 \quad \textit{and } \beta_0 = \frac{1}{q}, \tag{38}$$

where $G_{n,q} = G_{n,q}(0,0)$ is the n-th q-Genocchi number and can be obtained from the generating function $\frac{2t}{e_q(t)+1} = \sum_{n=0}^{\infty} G_{n,q}\frac{t^n}{[n]_q!}$.

Theorem 10. *The following linear homogeneous recurrence relation holds for the class of 2D q-Genocchi polynomials for every $n \geq 1$*

$$G_{n,q}(qx,y) = q^n(x+\frac{y-1}{q^2})G_{n-1,q}(x,y)+ \tag{39}$$
$$\frac{1}{2\,[n]_q}\sum_{k=1}^{n-1}\begin{bmatrix} n \\ k-1 \end{bmatrix}_q q^{k-2}G_{k-1,q}(x,y)(G_{n-k+1,q}+\frac{(q-1)y}{[n-k+1]_q}\sum_{l=0}^{n-k}\begin{bmatrix} n-k \\ l \end{bmatrix}_q G_{l,q}).$$

Theorem 11. *The following partial q-difference equations hold for the polynomials in the class of 2D q-Genocchi*

$$\left[(xq^n+\frac{y-1}{q})D_{q,x}+\sum_{k=2}^{n}\frac{q^{n-k-1}}{2[k]_q!}(G_{k,q}+\frac{(q-1)y}{[k]_q}\sum_{l=0}^{k-1}\begin{bmatrix} k-1 \\ l \end{bmatrix}_q G_{l,q})D_{q,x}^k\right] \tag{40}$$
$$\times\, G_{n,q}(x,y) - [n]_q G_{n,q}(qx,y) = 0.$$

$$(xq^n + \frac{y-q}{q^2})D_{q,y}G_{n,q}(x, \frac{y}{q}) + \sum_{k=2}^{n} \frac{q^{n-k-1}}{2[k]_q!}(G_{k,q} + \frac{(q-1)y}{q^k[k]_q} \sum_{l=0}^{k-1} \left[\begin{array}{c} k-1 \\ l \end{array} \right]_q G_{l,q})$$

(41)

$$\times D^k_{q,y}G_{n,q}(x, \frac{y}{q^k}) - [n]_q G_{n,q}(qx, y) = 0.$$

References

[1] Eini Keleshteri, M. & Mahmudov, N. I. (2015). A study on q-Appell polynomials from determinantal point of view. *Appl. Math. Comput.* 260 , 351369.

[2] Bretti, G. Natalini, P. & Ricci, P. E. (2002). Generalizations of the Bernoulli and Appell polynomials. *Abstract and Applied Analysis* , no. 2004, 7, 613623.

[3] Mahmudov, N. I. (2014). Difference equations of q-Appell polynomials. *Appl. Math. Comput.* 245, 539543.

[4] L. Carlitz, q-Bernoulli numbers and polynomials. (1948). *Duke Math. J.* 15, 987-1000.

[5] N.I. Mahmudov, On a class of q-Bernoulli and q-Euler polynomials. (2013). *Adv. Difference Equ.* , 2013, 108, 11 pp.

Mathematics Department, Eastern Mediterranean University, Famagusta, North Cyprus, via Mersin 10, Turkey

E-mail address: marzieh.eini@emu.edu.tr, nazim.mahmudov@emu.edu.tr